

Neutrino Masses in Theories with Dynamical Electroweak Symmetry Breaking

Thomas Appelquist^{a *}, Robert Shrock^{b †}

a Physics Department, Sloane Laboratory

Yale University 06520

New Haven, CT

(b) C. N. Yang Institute for Theoretical Physics

State University of New York

Stony Brook, N. Y. 11794

We address the problem of accounting for light neutrino masses in theories with dynamical electroweak symmetry breaking. We discuss this in the context of a class of (extended) technicolor (ETC) models and analyze the full set of Dirac and Majorana masses that arise in such theories. As a possible solution, we propose a combination of suppressed Dirac masses and a seesaw involving dynamically generated $|\Delta L| = 2$ condensates of standard-model singlet, ETC-nonsinglet fermions. We show how this can be realized in an explicit ETC model. An important feature of this proposal is that, because of the suppression of Dirac neutrino mass terms, a seesaw yielding realistic neutrino masses does not require superheavy Majorana masses; indeed, these Majorana masses are typically much smaller than the largest ETC scale.

14.60.PQ, 12.60.Nz, 14.60.St

*email: thomas.appelquist@yale.edu

†email: robert.shrock@sunysb.edu

I. INTRODUCTION

An understanding of the fermion mass spectrum remains an intriguing challenge for particle physics. The standard model (SM) accomodates quark and charged lepton masses by the mechanism of Yukawa couplings to a postulated Higgs boson, but this does not provide insight into these masses, especially since it requires small dimensionless Yukawa couplings for all of the charged fermions except the top quark, ranging down to $10^{-6} - 10^{-5}$ for the electron and u and d quarks. The standard model has zero neutrino masses, and hence must be modified to take account of the increasingly strong evidence for the very small but non-zero neutrino masses and significant lepton mixing from solar and atmospheric data [1,2], consistent with the K2K accelerator neutrino experiment [3].

Since masses for the quarks, charged leptons, and observed neutrinos break the chiral gauge symmetry of the standard model, an explanation of these masses necessarily involves a model for electroweak symmetry breaking (EWSB). One possibility is dynamical electroweak symmetry breaking driven by a strongly coupled gauge interaction, associated with an exact gauge symmetry, denoted generically as technicolor (TC) [4]- [10]. The EWSB arises from the condensation of technifermion bilinears. The generation of realistic masses for the charged leptons and u , d , s , c , and b quarks seems attainable in this framework, via extended technicolor, in particular with slowly running ("walking") technicolor. Although additional ingredients are very likely necessary to explain the large top-quark mass, we explore here the possibility that an ETC model of the above type can yield a plausible explanation for small neutrino masses. This is a significant challenge for dynamical EWSB models. As conventionally formulated, these theories have no very large mass scale analogous to the grand unification scale M_{GUT} that enters in the seesaw mechanism [11] yielding a Majorana mass $m_\nu \sim m_D^2/m_R$, where m_D is a Dirac mass and $m_R \sim M_{GUT}$ is the mass characterizing electroweak-singlet neutrinos.

Although some previous attempts have been made to study this problem [5,7,9], it is important to reconsider it in light of later theoretical and experimental developments. Refs. [5,9] did not include Majorana neutrino mass terms and instead explored a suppression mechanism for Dirac neutrino masses. This approach does not, however, yield enough suppression to agree with current experiments. Here we give a general treatment including both Dirac and Majorana mass terms. We show how ETC theories dynamically produce such Majorana mass terms and associated condensates, violating lepton number L as $|\Delta L| = 2$ [12]. We propose, as a possible solution for how to get light neutrino masses, a combination of naturally suppressed Dirac masses and a seesaw involving the dynamically generated Majorana mass terms. We show how this proposal can be realized in an explicit ETC model.

II. NEUTRINO MASS TERMS IN EXTENDED TECHNICOLOR THEORIES

We first present a general discussion taking the technicolor gauge group to be $SU(N_{TC})$. The set of technifermions includes, as a subset, one family, viz., $Q_L = \begin{pmatrix} U \\ D \end{pmatrix}_L$, $L_{TC,L} = \begin{pmatrix} N \\ E \end{pmatrix}_L$, U_R , D_R , N_R , E_R transforming according to the fundamental representation of $SU(N_{TC})$ and

the usual representations of $G_{SM} = \text{SU}(3) \times \text{SU}(2)_L \times \text{U}(1)_Y$ (color and TC indices are usually suppressed). To satisfy constraints from flavor-changing neutral-current processes, the ETC vector bosons, which can mediate generation-changing transitions, must have large masses. We envision that these arise from self-breaking of the ETC gauge symmetry, which requires that ETC be a strongly coupled, chiral gauge theory. The self-breaking occurs in stages, for example at the three stages $\Lambda_1 \sim 10^3$ TeV, $\Lambda_2 \sim 50$ TeV, and $\Lambda_3 \sim 3$ TeV, corresponding to the $N_{gen} = 3$ standard-model fermion generations. Then $N_{ETC} = N_{TC} + N_{gen}$.

A particularly attractive choice for the technicolor group, used in the explicit model to be studied here, is $\text{SU}(2)_{TC}$, which has the appeal that it minimizes the TC contributions to the S parameter [13] and can yield walking behavior, allowing for realistically large quark and charged lepton masses. With $N_{gen} = 3$, the choice $N_{TC} = 2$ corresponds to $N_{ETC} = 5$. With $N_f = 8$ vectorially coupled technifermions in the fundamental representation, studies suggest that this $\text{SU}(2)_{TC}$ theory could have an (approximate) infrared fixed point (IRFP) in the confining phase with spontaneous chiral symmetry breaking but near to the phase transition (as a function of N_f for fixed N_{TC}) beyond which the theory would go over into a nonabelian Coulomb phase [14,15]. This approximate IRFP provides the walking behavior, enhancing the technifermion condensates that control the quark and charged lepton masses. The walking can also enhance the masses of pseudo-Nambu-Goldstone bosons, but further ingredients are likely needed to ensure the absence of some massless Nambu-Goldstone bosons.

A rough estimate of the quark and charged lepton masses can be made by considering a one-loop diagram in which a fermion f_a emits a virtual ETC gauge boson, going to a virtual technifermion F which reabsorbs the gauge boson, producing the mass term $m_{f_a} \bar{f}_{a,L} f_{a,R} + h.c.$ with

$$m_{f_a} \sim \frac{g_{ETC}^2 \eta_a N_{TC} \Lambda_a^3}{4\pi^2 M_a^2} \quad (1)$$

where $M_a \sim g_{ETC} \Lambda_a$ is the mass of the ETC gauge bosons that gain mass at scale Λ_a and g_{ETC} is the running ETC gauge coupling evaluated at this scale. In eq. (1) η_a is a possible enhancement factor incorporating walking, which can be as large as Λ_a/f_F [6,16], where f_F is the technicolor pseudoscalar decay constant (for our purposes we can take $f_L \simeq f_Q \equiv f_F$). We recall that Λ_{TC} is determined by using the relation $m_W^2 = (g^2/4)(N_c f_Q^2 + f_L^2) \simeq (g^2/4)(N_c + 1)f_F^2$, which gives $f_F \simeq 130$ GeV. In QCD, $f_\pi = 93$ MeV and $\Lambda_{QCD} \sim 170$ MeV, so that $\Lambda_{QCD}/f_\pi \sim 2$; using this as a guide to technicolor, we infer $\Lambda_{TC} \sim 260$ MeV.

Technicolor models in general also have a set of electroweak-singlet neutrinos, $\chi_R = (\chi_1, \dots, \chi_{n_s})_R$ [17], some technicolored and some techni-singlets, in addition to the left-handed, weak-isospin-doublet neutrinos and technineutrinos. The contributions to the total neutrino mass matrix, generated by condensates arising at the TC and ETC scales, are then of three types: (i) left-handed Majorana, (ii) Dirac, and (iii) right-handed Majorana. The left-handed Majorana mass terms, which violate L by two units, take the form

$$\sum_{i,j=1}^{N_{ETC}} [n_{iL}^T C(M_L)_{ij} n_{jL}] + h.c. \quad (2)$$

where $n_L = (\{\nu_\ell\}, \{N\})_L$ includes the electroweak-doublet left-handed neutrinos for $i, j = 1, 2, 3$ and technineutrinos for $i, j = 4, \dots, N_{ETC}$; and $C = i\gamma_2\gamma_0$. Left-handed Majorana masses violate the electroweak gauge symmetry, and, for technineutrinos, also the TC symmetry, which is exact. Thus, $(M_L)_{ij} = 0$ for i or $j = 4, \dots, N_{ETC}$. The Dirac mass terms take the form

$$\sum_{a=1}^{N_{ETC}} \sum_{s=1}^{n_s} \bar{n}_{aL} (M_D)_{as} \chi_{sR} + h.c. \quad (3)$$

Finally, the Majorana bilinears with SM-singlet neutrinos are

$$\sum_{s,s'=1}^{n_s} \chi_{sR}^T C (M_R)_{ss'} \chi_{s'R} , \quad (4)$$

In (3) and (4) $(M_D)_{as} = 0$ and $(M_R)_{ss'} = 0$ for technicolor-noninvariant entries.

The full neutrino mass term is then

$$-\mathcal{L}_m = \frac{1}{2} (\bar{n}_L \bar{\chi}_L^c) \begin{pmatrix} M_L & M_D \\ (M_D)^T & M_R \end{pmatrix} \begin{pmatrix} n_R^c \\ \chi_R \end{pmatrix} + h.c. \quad (5)$$

Since $(M_L)^T = M_L$ and $(M_R)^T = M_R$, the full $(N_{ETC} + n_s) \times (N_{ETC} + n_s)$ neutrino mass matrix M in (5) is complex symmetric and can be diagonalized by a unitary transformation $U_\nu^\dagger$ as $M_{diag.} = U_\nu^\dagger M (U_\nu^\dagger)^T$. This yields the neutrino masses and transformation U_ν relating the group eigenstates $\nu_L = (\bar{n}, \bar{\chi}^c)_L^T$ and the corresponding mass eigenstates $\nu_{m,L}$, according to $\nu_{j,L} = \sum_{k=1}^{N_{ETC}+n_s} (U_\nu)_{jk} \nu_{k,m,L}$, $1 \leq j \leq N_{ETC} + n_s$ (the elements $(U_\nu)_{jk}$ connecting technisingle and technicolored neutrinos vanish identically). The lepton mixing matrix for the observed neutrinos [18] $\nu_{\ell,L} = U \nu_{m,L}$ is then given by

$$U_{ik} = \sum_{j=1}^3 (U_{\ell,L})_{ij} (U_\nu)_{jk} , \quad 1 \leq i \leq 3, \quad 1 \leq k \leq N_{ETC} + n_s \quad (6)$$

where $U_{1k} \equiv U_{ek}$, etc., and where the diagonalization of the charged lepton mass matrix is carried out by the bi-unitary transformation $M_{\ell,diag.} = U_{\ell,L} M_\ell U_{\ell,R}^\dagger$.

III. SPECIFIC EXTENDED TECHNICOLOR MODEL

We next present an analysis of a specific ETC model based on the gauge group $G = SU(5)_{ETC} \times SU(2)_{HC} \times G_{SM}$. One additional gauge interaction, $SU(2)_{HC}$, where HC denotes hypercolor, has been introduced along with $SU(5)_{ETC}$ and G_{SM} . Both the $SU(2)_{HC}$ and $SU(5)_{ETC}$ interactions become strong, triggering a sequential breaking pattern. The fermion content of this model is listed below, where the numbers indicate the representations under $SU(5)_{ETC} \times SU(2)_{HC} \times SU(3)_c \times SU(2)_L$ and the subscript gives the weak hypercharge:

$$\begin{aligned}
& (5, 1, 3, 2)_{1/3, L} , \quad (5, 1, 3, 1)_{4/3, R} , \quad (5, 1, 3, 1)_{-2/3, R} \\
& (5, 1, 1, 2)_{-1, L} , \quad (5, 1, 1, 1)_{-2, R} , \quad (\overline{10}, 1, 1, 1)_{0, R} , \\
& (10, 2, 1, 1)_{0, R} .
\end{aligned} \tag{7}$$

Thus the fermions include quarks and techniquarks in the representations $(5, 1, 3, 2)_{1/3, L}$, $(5, 1, 3, 1)_{4/3, R}$, and $(5, 1, 3, 1)_{-2/3, R}$, left-handed charged leptons and neutrinos and technileptons in $(5, 1, 1, 2)_{-1, L}$, and right-handed charged leptons and technileptons in $(5, 1, 1, 1)_{-2, R}$, together with SM-singlet fermions $\psi_{ij, R}$ in the antisymmetric tensor representation $(\overline{10}, 1, 1, 1)_{0, R}$. The unusual assignment of the SM singlets makes the $SU(5)_{ETC}$ gauge theory chiral. Finally, in order to render the theory anomaly-free and to provide interactions to help trigger the symmetry breaking, one adds the hypercolored fields in the $(10, 2, 1, 1)_{0, R}$, denoted $\zeta_R^{ij, \alpha}$, where ij and α are ETC and HC indices. Thus, $n_s = 30$. We label the ETC gauge bosons as $(V_j^i)_\mu$, $1 \leq i, j \leq 5$. To fix the convention for the lepton number assigned to $\psi_{ij, R}$, we take it to be $L = 1$ in order that Dirac terms $\bar{n}_{i, L} \psi_{jk, R}$ conserve lepton number. The lepton number assigned to the $\zeta_R^{ij, \alpha}$ fields is also a convention; since they have no Dirac terms with observed neutrinos, we leave it arbitrary. We write $\chi_R = (\psi, \zeta)_R$.

Each of the nonabelian factor groups in G is asymptotically free. There are no bilinear fermion operators invariant under G and hence there are no bare fermion mass terms. The $SU(2)_{HC}$ and $U(1)_{HC}$ interactions and the $SU(2)_{TC}$ subsector of $SU(5)_{TC}$ are vectorial. This model has some features in common with the ETC model, denoted AT94, of [9], but has different gauge groups and fermion content.

We next analyze the stages of symmetry breaking. We envision that at $E \sim \Lambda_1 \sim 10^3$ TeV, α_{ETC} is sufficiently large to produce condensation in the attractive channel $(\overline{10}, 1, 1, 1)_{0, R} \times (\overline{10}, 1, 1, 1)_{0, R} \rightarrow (5, 1, 1, 1)_0$, breaking $SU(5)_{ETC} \rightarrow SU(4)_{ETC}$. In the most attractive channel (MAC) analysis this is a highly attractive channel, with $\Delta C_2 = 24/5$, although it is not the MAC itself. (The MAC is $(\overline{10}, 1, 1, 1)_{0, R} \times (10, 2, 1, 1)_{0, R} \rightarrow (1, 2, 1, 1)$, with $\Delta C_2 = 36/5$; this is undesired since it would break $SU(2)_{HC}$.) The desired condensation channel is nearly as strong and is just as probable within the uncertainties of MAC analyses. With no loss of generality, we take the breaking direction in $SU(5)_{ETC}$ as $i = 1$; this entails the separation of the first generation of quarks and leptons from the components of $SU(5)_{ETC}$ fields with indices lying in the set $\{2, 3, 4, 5\}$. With respect to the unbroken $SU(4)_{ETC}$, we have the decomposition $(\overline{10}, 1, 1, 1)_{0, R} = (\bar{4}, 1, 1, 1)_{0, R} + (\bar{6}, 1, 1, 1)_{0, R}$. We denote the $(\bar{4}, 1, 1, 1)_{0, R}$ and antisymmetric tensor representation $(\bar{6}, 1, 1, 1)_{0, R}$ as $\alpha_{1iR} \equiv \psi_{1i, R}$ for $2 \leq i \leq 5$ and $\xi_{ij, R} \equiv \psi_{ij, R}$ for $2 \leq i, j \leq 5$. The associated $SU(5)_{ETC}$ -breaking, $SU(4)_{ETC}$ -invariant condensate is then

$$\langle \epsilon_{1ijkl} \zeta_R^{ijT} C \xi_R^{k\ell} \rangle = 4 \langle \xi_R^{23T} C \xi_R^{45} - \xi_R^{24T} C \xi_R^{35} + \xi_R^{25T} C \xi_R^{34} \rangle . \tag{8}$$

This condensate and the resultant dynamical Majorana mass terms for the six components of ξ in eq. (8) violate total lepton number as $|\Delta L| = 2$. The dynamical formation of Majorana mass terms and violation of total lepton number is an important feature of these models, providing a necessary ingredient for a (dynamical) seesaw mechanism [20].

At lower scales, depending on relative strengths of couplings, different symmetry-breaking sequences occur. One plausible sequence, denoted G_a , is as follows: at $\Lambda_2 \sim 10^2$ TeV, $SU(4)_{ETC}$ and $SU(2)_{HC}$ couplings are sufficiently large to lead together to the condensation $(4, 2, 1, 1)_{0,R} \times (6, 2, 1, 1)_{0,R} \rightarrow (\bar{4}, 1, 1, 1)$, breaking $SU(4)_{ETC} \rightarrow SU(3)_{ETC}$ [9]. This condensate is

$$\langle \epsilon_{\alpha\beta} \epsilon_{i2jk\ell} \zeta_R^{ij,\alpha} {}^T C \zeta_R^{k\ell,\beta} \rangle = 4 \langle \epsilon_{\alpha\beta} (\zeta_R^{13,\alpha} {}^T C \zeta_R^{45,\beta} - \zeta_R^{14,\alpha} {}^T C \zeta_R^{35,\beta} + \zeta_R^{15,\alpha} {}^T C \zeta_R^{34,\beta}) \rangle, \quad (9)$$

and the twelve $\zeta_R^{ij,\alpha}$ fields in this condensate gain masses $\sim \Lambda_2$. Both the $SU(4)_{ETC}$ and $SU(2)_{HC}$ interactions are strongly attractive in this channel, together making the channel an example of the big-MAC of Ref. [9]. The fact that the neutrino-like fields $\alpha_{1i,R}$ transform as a $\bar{4}$ of $SU(4)_{ETC}$, while the left-handed neutrinos and technineutrinos transform as a 4, will lead to a strong suppression of relevant entries in the Dirac submatrix M_D [5,9].

In the G_a symmetry-breaking sequence, at the lowest ETC scale, $\Lambda_3 \sim 3$ TeV, the $(3, 2, 1, 1)_{0,R}$, $\zeta_R^{2j,\alpha}$, $j = 3, 4, 5$, from the $(6, 2, 1, 1)_{0,R}$ is assumed to condense as $(3, 2, 1, 1)_{0,R} \times (3, 2, 1, 1)_{0,R} \rightarrow (\bar{3}, 1, 1, 1)$, breaking $SU(3)_{ETC} \rightarrow SU(2)_{TC}$ [9]. The condensate is $\langle \epsilon_{\alpha\beta} \zeta_R^{24,\alpha} {}^T C \zeta_R^{25,\beta} \rangle$. This breaking again involves the combination of attractive ETC and HC interactions [9]. Further, we expect that at a scale $\sim \Lambda_3$ the HC interaction produces the condensate $\langle \epsilon_{\alpha\beta} \zeta_R^{12,\alpha} {}^T C \zeta_R^{23,\beta} \rangle$. Thus, just as the six $\xi_{ij,R}$ condense out of the theory at energies below Λ_1 , all of the 20 fields $\zeta^{ij,\alpha}$ in the $(10, 2, 1, 1)_{0,R}$ have condensed out of the effective theory at energies below Λ_3 . Since one may assign lepton number zero to $\zeta_R^{ij,\alpha}$, condensates of the form $\langle \zeta_R^T C \zeta_R \rangle$ do not necessarily violate total lepton number in this model.

A different sequence of condensations, denoted G_b , can occur if the $SU(2)_{HC}$ coupling is somewhat smaller. At a scale $\Lambda_{BHC} \lesssim \Lambda_1$ (BHC = broken HC), the $SU(4)_{ETC}$ interaction produces a condensation in the channel $(6, 2, 1, 1)_{0,R} \times (6, 2, 1, 1)_{0,R} \rightarrow (1, 3, 1, 1)_0$. With respect to ETC, this channel has $\Delta C_2 = 5$ and is hence slightly more attractive than the initial condensation (8) with $\Delta C_2 = 24/5$, but it can occur at the somewhat lower scale Λ_{BHC} because it is repulsive with respect to hypercolor. With no loss of generality, one can orient $SU(2)_{HC}$ axes so that the condensate is

$$\langle \epsilon_{1ijk\ell} \zeta_R^{ij,1} {}^T C \zeta_R^{k\ell,2} \rangle + (1 \leftrightarrow 2). \quad (10)$$

Since this is an adjoint representation of hypercolor, it breaks $SU(2)_{HC} \rightarrow U(1)_{HC}$. We let $\alpha = 1, 2$ correspond to $Q_{HC} = \pm 1$ under the $U(1)_{HC}$. This gives dynamical masses $\sim \Lambda_{BHC}$ to the twelve $\zeta_R^{ij,\alpha}$ fields involved.

At a lower scale, Λ_{23} , in the G_b sequence, we envision that a combination of the $SU(4)_{ETC}$ and $U(1)_{HC}$ attractive interactions produces the condensation $4 \times 4 \rightarrow 6$ with condensate $\langle \epsilon_{\alpha\beta} \zeta_R^{12,\alpha} {}^T C \zeta_R^{13,\beta} \rangle$, which then breaks $SU(4)_{ETC} \rightarrow SU(2)_{ETC}$ and is $U(1)_{HC}$ -invariant. Thus, the sequence G_b has only two ETC breaking scales, Λ_1 and Λ_{23} ; additional ingredients are needed to obtain the requisite range of SM fermion masses. Here we take $\Lambda_{23} \sim 10$ TeV. Although there is a residual $U(1)_{HC}$ gauge interaction in these models, its effects are shielded since it does not couple directly to SM particles. Finally, for both G_a and G_b , at the still lower scale $\Lambda_{TC} \sim f_F$, technifermion condensation takes place, breaking $SU(2)_L \times U(1)_Y \rightarrow U(1)_{em}$.

IV. CALCULATIONS AND RESULTS

The mass matrix M of neutrino-like (colorless and electrically neutral) states in Eq. (5) has $N_{ETC} = 5$ and $n_s = 30$. Since the hypercolored fields do not form bilinear condensates and resultant mass terms with hypercolor singlets, M is block-diagonal, comprised of a 15×15 block M_{HCS} involving hypercolor-singlet neutrinos and a 20×20 block M_{HC} involving the hypercolored fermions, M_{HC} . The entries in the matrix M arise as the high-energy physics is integrated out at each stage of condensation from Λ_1 down to Λ_{TC} . Composite operators of various dimension are formed, with bilinear condensation then leading to the masses. The nonzero entries of M arise in two different ways: (i) directly, as dynamical masses associated with various condensates, and (ii) via loop diagrams involving dynamical mass insertions on internal fermion lines and, in most cases, also mixings among ETC gauge bosons on internal lines. Since the ETC gauge boson mixing arises at the level of one or more loops, most graphs for nonzero type-(ii) elements of M arise at the level of at least two-loop diagrams. The different origins for the elements of M give rise to quite different magnitudes for these elements; in particular, there is substantial suppression of most type-(ii) entries. This suppression is not primarily due to the ETC gauge couplings, which are strong, but to the fact that the diagrams involve ratios of small scales such as Λ_{TC} and lower ETC scales to larger scales such as Λ_1 . The 20×20 matrix M_{HC} involving the $(10, 2, 1, 1)_{0,R}$ fermions contains dynamical fermion mass entries resulting from the hypercolor condensates and has $\text{Tr}(M_{HC}) = 0$.

The matrix of primary interest, M_{HCS} , is given by the operator product

$$-\mathcal{L}_{HCS} = \frac{1}{2}(\bar{n}_L, \bar{\alpha}_L^c, \bar{\xi}_L^c) \begin{pmatrix} M_L & (M_D)_{\bar{n}\alpha} & (M_D)_{\bar{n}\xi} \\ (M_D)_{\bar{n}\alpha}^T & (M_R)_{\alpha\alpha} & (M_R)_{\alpha\xi} \\ (M_D)_{\bar{n}\xi}^T & (M_R)_{\alpha\xi}^T & (M_R)_{\xi\xi} \end{pmatrix} \begin{pmatrix} n_R^c \\ \alpha_R \\ \xi_R \end{pmatrix} + h.c. \quad (11)$$

The five-component n_R^c , the four-component α_R , and the six-component ξ_R each contain TC singlets as well as nonsinglets. One of the two Dirac submatrices is

$$(M_D)_{\bar{n}\alpha} = \begin{pmatrix} b_{12} & b_{13} & 0 & 0 \\ b_{22} & b_{23} & 0 & 0 \\ b_{32} & b_{33} & 0 & 0 \\ 0 & 0 & 0 & c_1 \\ 0 & 0 & -c_1 & 0 \end{pmatrix} \quad (12)$$

The vanishing entries are zero because of exact technicolor gauge invariance. The entry c_1 represents a dynamical mass directly generated by technicolor interactions corresponding to $\sum_{i,j=4,5} \epsilon^{ij} \bar{n}_{i,L} \alpha_{1j,R}$, so that $|c_1| \sim \Lambda_{TC}$. Note that this involves the antisymmetric, ϵ^{ij} , rather than the δ_j^i , contraction of $\text{SU}(2)_{TC}$ indices and thus makes crucial use of the fact that the technicolor group is $\text{SU}(2)$ rather than $\text{SU}(N)$ with $N \geq 3$.




FIG. 1. Graphs generating $\bar{n}_{i,L} b_{ij} \alpha_{1j,R}$ for $i = 1, 2, 3$ and $j = 2, 3$, assuming that the indicated mixings of ETC gauge bosons occur.




FIG. 2. One-loop graph contributing to the gauge boson mixing $V_3^4 \leftrightarrow V_5^2$. The graph with indices 4 and 5 interchanged on the internal ζ lines also contributes.

In Fig. 1 we show graphs that could yield the b_{ij} 's. Here the $\times$ on the fermion line represents the dynamical mass corresponding to a technicolor condensate. Each graph requires nondiagonal insertions on the internal ETC gauge bosons lines. We find that the requisite ETC gauge boson mixings occur to leading (one-loop) order in the G_a sequence for (i) b_{13} , which involves $V_3^4 \leftrightarrow V_5^1$ and $V_3^5 \leftrightarrow V_4^1$, and (ii) b_{22} , which involves $V_2^4 \leftrightarrow V_5^2$; and in the G_b sequence for b_{23} and b_{32} , which involve $V_3^4 \leftrightarrow V_5^2$ and $V_3^5 \leftrightarrow V_4^2$. For example, for G_b we show in Fig. 2 the one-loop graphs contributing to $V_3^4 \leftrightarrow V_5^2$. In each respective case, G_a and G_b , the other b_{ij} 's are produced by higher-loop diagrams. For example, starting from Fig. 1 for b_{23} in case G_b , one can construct diagrams in which the incoming $\alpha_{13,R}$ or the virtual $\alpha_{14,R}$ or n_L^5 emits a virtual V_k^k ETC gauge boson with $k \in \{1, 2, 3\}$ which, via mixing, becomes V_2^i , which is then absorbed by the n_L^2 to yield an outgoing n_L^i , $i = 1, 3$. Other similar graphs involving a triple ETC gauge-boson vertex along with mixing also contribute in this way. These generate b_{13} and b_{33} at a level suppressed relative to b_{23} . The $V_k^k \rightarrow V_2^i$ mixing arises

generically from loop graphs in which at least one internal fermion line is a standard-model quark or charged lepton with a mass insertion that is nondiagonal in generation, incorporating the mixing of the weak eigenstates of these fermions to form mass eigenstates. The entries b_{12} and b_{22} are generated in a similar way.

We next estimate the leading b_{ij} entries. For either breaking sequence, we denote the ETC gauge boson 2-point function as

$${}^k\Pi_j^i(q)_{\mu\lambda} = \int \frac{d^4x}{(2\pi)^4} e^{iq \cdot x} \langle T [(V_n^k)_\mu(x/2)(V_j^i)_\lambda(-x/2)] \rangle_0 . \quad (13)$$

After some manipulations (and Wick rotation), the graph in Fig. 1 yields

$$g_{ETC}^2 [\bar{n}_{i,L}(p) \gamma_\mu \gamma_\lambda \alpha_{1j,R}(p)] \int \frac{d^4k}{(2\pi)^4} \frac{k^2 \Sigma_{TC}(k) [{}_5^i\Pi_j^4((p-k)^2)]^{\mu\lambda}}{(k^2 + \Sigma_{TC}(k)^2)^2 [(p-k)^2 + M_j^2][(p-k)^2 + M_i^2]} , \quad (14)$$

where $\Sigma_{TC}(k)$ is the dynamical technicolor mass associated with the transition $\alpha_{14,R} \rightarrow n_L^5$. This mass has the behavior $\Sigma_{TC}(k) \sim \Lambda_{TC}$ for $k^2 \ll \Lambda_{TC}^2$, while for $k^2 \gg \Lambda_{TC}^2$, (i) $\Sigma_{TC}(k) \sim \Lambda_{TC}^2/k$ for a walking theory [6], (ii) $\Sigma_{TC}(k) \sim \Lambda_{TC}^3/k^2$ in a QCD-like theory. Hence, we need ${}_n^k\Pi_j^i((p-k)^2)_{\mu\lambda}$ only for $(p-k)^2/\Lambda_1^2 \ll 1$, since the loop momenta in Fig. 1 are cut off far below Λ_1 (at Λ_3 for G_a or Λ_{23} for G_b). In eq. (14), M_j denotes the mass of the ETC gauge boson that picks up mass at Λ_j .

In the sequence G_b , for $q^2 \ll \Lambda_1^2$, we estimate

$$[{}_5^2\Pi_3^4(q)]_{\mu\lambda} \sim [{}_4^2\Pi_3^5(q)]_{\mu\lambda} \sim \frac{g_{ETC}^2 \Lambda_{TC}^2}{(2\pi^2)} g_{\mu\lambda} . \quad (15)$$

where we have assumed a walking behavior of the TC theory up to Λ_{23} . For $i, j = 2, 3$ and $3, 2$, adding the other graph with $4 \leftrightarrow 5$ in Fig. 1, we find

$$|b_{23}| = |b_{32}| \sim \frac{g_{ETC}^4 \Lambda_{TC}^4 \Lambda_{23}}{2\pi^4 M_{23}^4} \sim \frac{\Lambda_{TC}^4}{2\pi^4 \Lambda_{23}^3} \quad \text{for } G_b , \quad (16)$$

where we have again assumed the above walking TC behavior. For sequence G_a , we estimate, using similar methods,

$$|b_{13}| \sim \frac{\Lambda_{TC}^2 \Lambda_3}{2\pi^4 \Lambda_1^2} , \quad |b_{22}| \sim \frac{\Lambda_{TC}^2 \Lambda_3}{2\pi^4 \Lambda_2^5} \quad \text{for } G_a . \quad (17)$$

With the numerical inputs given above, we get $|b_{23}| = |b_{32}| \sim O(1)$ KeV for G_b and $|b_{13}| \sim O(1)$ KeV and $|b_{22}| \sim O(10)$ eV for G_a . Because the ETC and TC theories are strongly coupled, these estimates based on perturbative expansions in powers of α_{ETC} involve an obvious uncertainty. For each case, the other b_{ij} 's are generated at smaller levels. These calculations show how this aspect - suppressed Dirac neutrino masses - of our proposal are realized in an explicit model. While the specific results for the various b_{ij} are dependent on the model and symmetry breaking pattern, one can infer that this type of suppression can

be achieved in a general class of ETC models where Dirac mass terms are generated in a similar manner.

The second Dirac submatrix in eq. (11) is

$$(M_D)_{\bar{n}\xi} = \begin{pmatrix} d_{1,23} & d_{1,45} & 0 & 0 & 0 & 0 \\ d_{2,23} & d_{2,45} & 0 & 0 & 0 & 0 \\ d_{3,23} & d_{3,45} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & c_2 & 0 & c_3 \\ 0 & 0 & -c_2 & 0 & -c_3 & 0 \end{pmatrix} \quad (18)$$

Again, the zeros are exact and follow from technicolor invariance. Because the ξ fields decouple from the theory at scales below Λ_1 , the nonzero elements of $(M_D)_{\bar{n}\xi}$ arise indirectly, via loop diagrams and are highly suppressed. These elements of $(M_D)_{\bar{n}\xi}$ have only a small effect on the neutrino eigenvalues because in the characteristic polynomial $P(x)$ they occur as corrections to much larger terms involving Λ_1 .




FIG. 3. Graphs for $\alpha_{12,R}^T C r_{23} \alpha_{13,R}$ in case G_b .

In M_R the 6×6 submatrix $(M_R)_{\xi\xi}$ has six nonzero entries that are dynamical mass terms of order Λ_1 arising directly from the condensate (8). These are important since they are $|\Delta L| = 2$ operators, and they, in turn, induce the $(M_R)_{\alpha\alpha}$ Majorana mass terms which play a central role in the seesaw. Thus the $(M_R)_{\xi\xi}$ entries are the underlying seed for the Majorana mass terms involving the observed neutrinos. Note that $\text{Tr}(M_R) = 0$.

The submatrix $(M_R)_{\alpha\alpha}$ has the form

$$(M_R)_{\alpha\alpha} = \begin{pmatrix} r_{22} & r_{23} & 0 & 0 \\ r_{23} & r_{33} & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (19)$$

As before, the zeros are exact and are due to technicolor invariance. If the 2×2 r_{ij} submatrix has maximal rank, this can provide a seesaw which, in conjunction with the suppression of the Dirac entries b_{ij} discussed above, can yield adequate suppression of neutrino masses. The submatrix r_{ij} , $2 \leq i, j \leq 3$, produces this seesaw because $\alpha_{12,R}$ and $\alpha_{13,R}$ are the electroweak-singlet techni-singlet neutrinos that remain as part of the low-energy effective field theory at and below the electroweak scale.

Consider the sequence G_b . In Fig. 3 we show graphs contributing to r_{23} for this case. These depend on the $V_1^4 \leftrightarrow V_5^1$ ETC gauge mixing produced by the graphs in Fig. 4. From these we calculate

$$r_{23} \sim \frac{\Lambda_{BHC}^2 \Lambda_{23}^2}{2\pi^4 \Lambda_1^3} \quad \text{for } G_b. \quad (20)$$

where here we have assumed a walking behavior of the ETC theory below Λ_{BHC} . The entries r_{22} and r_{33} are generated by higher-loop diagrams starting from the graphs in Fig. 3 for r_{23} in a manner similar to that whereby subdominant b_{ij} are generated starting from Fig. 1 for b_{23} and b_{32} . Numerically, with the above inputs, $|r_{23}| \sim O(0.1)$ GeV, with smaller values for r_{ii} , $i = 2, 3$. For sequence G_a we find that the r_{ij} entries are generated via higher-loop diagrams analogous to those for r_{22} and r_{33} in sequence G_b and hence are smaller than eq. (20). In the estimates to follow we concentrate on the sequence G_b since it yields a phenomenologically more successful seesaw, although this sequence has only two ETC breaking scales.




FIG. 4. One-loop graph for the ETC gauge boson mixing $V_1^4 \leftrightarrow V_5^1$ in case G_b . The graph with indices 4 and 5 interchanged on the internal ζ lines also contributes.

In the 4×6 submatrix $(M_R)_{\alpha\xi}$ the entries are either exactly zero by technicolor invariance or are nonzero but highly suppressed because the ξ fields decouple from the effective theory below Λ_1 . The nonzero entries do not have an important effect on the masses of neutrino-like states because of the way that they enter in the characteristic polynomial (similar to the elements of $(M_D)_{\bar{n}\xi}$).

We next summarize the above discussion from the viewpoint of effective field theory. At energy scales below Λ_{TC} , in either the breaking sequence G_a or G_b , the sector of neutrino-like states consists of the the techni-singlet components $i = 1, 2, 3$ of n_L^i and the techni-singlet components $\alpha_{1i,R}$, $i = 2, 3$; other fields have gained masses at higher scales and have been integrated out. The effective theory comprised of these degrees of freedom involves bilinear

(mass) operators along with a tower of higher-dimension operators. The mass operators are either of the Dirac type (the b_{ij} terms of eq. (11)) or of the Majorana type (the r_{ij} of eq. (19)). They form a 5×5 submatrix of M_{HCS} , and their magnitudes, which depend on the specific breaking sequence, are $\ll \Lambda_{TC}$.

Integrating out the $\alpha_{12,R}$ and $\alpha_{13,R}$ fields then yields the lowest-scale effective field theory, in which there are three light fermions, the n_L^i . The mass terms in this theory correspond to elements of M_L , and there are also higher-dimension operators involving the n_L^i . With respect to the mass terms, this procedure corresponds to a block diagonalization (“block-seesaw”) of the 5×5 submatrix of M_{HCS} , keeping only the light, M_L matrix. Its dominant terms arise in this manner; other, smaller entries are generated via higher-loop diagrams involving higher-dimension operators, for example induced by the exchange and mixing of ETC gauge bosons. The final step in the effective-field-theory approach is to diagonalize this 3×3 matrix, leading to the neutrino mass eigenvalues and mixing angles. Equivalently, one can think in terms of diagonalizing the full M_{HCS} -matrix in one fell swoop.

To be specific, we focus on the G_b sequence since it most clearly yields a seesaw. The largest M_L entry is $(M_L)_{23}$ (since $M_L = M_L^T$, we take $i \leq j$), and other, smaller terms arise from higher dimension operators. The electroweak-nonsinglet neutrinos are, to very good approximation, linear combinations of three mass eigenstates, of which the heaviest is ν_3 or ν_2 and has a mass

$$m_{\nu,max} \sim \frac{|b_{23}b_{32}|}{|r_{23}|} \sim \frac{\Lambda_{TC}^8 \Lambda_1^3}{2\pi^4 \Lambda_{23}^8 \Lambda_{BHC}^2} . \quad (21)$$

With the above-mentioned numerical values and $\Lambda_{BHC} \simeq 0.3\Lambda_1$, we find $m_{\nu,max} \simeq 0.05$ eV, consistent with experimental indications [2] based on a hierarchical spectrum, in which $m_{\nu,max} \simeq \sqrt{\Delta m_{32}^2}$. The model naturally yields large $\nu_\mu - \nu_\tau$ mixing because of the leading off-diagonal structure of the b_{ij} and r_{ij} with $ij = 23$ and 32 . The value of $|\Delta m_{32}^2|$ depends on details of the model but is on the low side of the experimental range. The lightest neutrino mass, $m(\nu_1)$, arises from the subdominant terms in M_L and is therefore predicted to be considerably smaller than $m(\nu_i)$, $i = 2, 3$. The group eigenstates involved in these (Majorana) mass eigenstates are $n_{i,R}^c$, $i = 1, 2, 3$ and $\alpha_{1j,R}$, $j = 2, 3$. This model thus exhibits our proposed explanation for light neutrino masses incorporating highly suppressed Dirac neutrino mass entries, $|\Delta L| = 2$ neutrino condensates and associated dynamical Majorana mass terms, and a resultant seesaw.

The model also yields the following mass eigenvalues and corresponding eigenvectors for the other neutrino-like states: (i) linear combinations (LC’s) of components of the six $\xi_{ij,R}$ with $2 \leq i, j \leq 5$ get masses $\sim \Lambda_1$; (ii) LC’s of the $\zeta_R^{ij,\alpha}$ with $2 \leq i, j \leq 5$ get masses $\sim \Lambda_{BHC}$; (iii) LC’s of the $\zeta^{1j,\alpha}$ with $j = 1, 2$ get masses $\sim \Lambda_{23}$; (iv) for technicolor nonsinglets, LC’s of the $\zeta^{1j,\alpha}$ with $j = 4, 5$ and LC’s of $n_{i,R}^c$ and $\alpha_{1i,R}$, with $i = 4, 5$ get masses $\sim \Lambda_{TC}$; (v) LC’s of $\alpha_{1i,R}$ with $i = 2, 3$ get masses $\sim r_{23}$. These masses are (nearly) Dirac.

Not only are the m_R entries responsible for the seesaw not superheavy masses; they are actually much smaller than the ETC scales Λ_i . A generic prediction of ETC models with the proposed seesaw is that some components of SM-singlet neutrino group eigenstates comprise dominant parts of mass eigenstates with masses given by the elements in M_R that

are involved in the seesaw (here, r_{23}). A condition to fit current limits on the emission of massive neutrinos, via lepton mixing, in particle decays would be that the $|U_{ek}|^2$, $|U_{\mu k}|^2 \lesssim 10^{-7}$ for $k > 3$ [21,22], which can be met while also maintaining sufficiently short lifetimes to satisfy astrophysical constraints.

V. CONCLUSIONS

In summary, we have given a general analysis of neutrino masses in the context of dynamical electroweak symmetry breaking theories, taking account of both Dirac and Majorana mass terms. We proposed a possible solution to the problem of obtaining light neutrino masses in this class of theories. This solution involves two main parts: (i) strong suppression of Dirac neutrino masses, and (ii) dynamical formation of bilinear Majorana neutrino condensates at ETC scales and resultant Majorana masses violating total lepton number as $|\Delta L| = 2$, and consequently a seesaw mechanism. We have shown how this proposal can be realized in an explicit ETC model. While further work is needed to obtain the detailed structural features needed to fit current indications for neutrino masses and lepton mixing, we believe that our proposal contains key ingredients for a solution to this problem in the context of theories with dynamical electroweak symmetry breaking. An important aspect of this suggestion is that it does not need any superheavy scale for a viable seesaw; indeed, the relevant Majorana masses may be much smaller than the highest ETC scale.

This research was partially supported by the grants DE-FG02-92ER-4074 (T.A.) and NSF-PHY-00-98527 (R.S.). T.A. thanks B. Dobrescu for helpful comments.

-
- [1] S. Fukuda, et al. Phys. Rev. Lett. **86** (2001) 5651, 5656; Phys. Lett. **B539** (2002) 179 (SuperK); Q. Ahmad et al., Phys. Rev. Lett. **87** (2001) 071301; *ibid.* **89** (2002) 011301, 011302 (SNO). Other data is from the Homestake, Kamiokande, GALLEX, and SAGE experiments. The optimal fit to this data involves ν_e oscillations into ν_μ and ν_τ with $\Delta m_{21}^2 \sim 5 \times 10^{-5} \text{ eV}^2$, where $\Delta m_{ij}^2 = m(\nu_i)^2 - m(\nu_j)^2$, and a relatively large associated mixing angle.
 - [2] Y. Fukuda et al., Phys. Lett. **B433** (1998) 9; Phys. Rev. Lett. **81** (1998) 1562; *ibid.*, **82** (1999) 2644; Phys. Lett. **B467** (1999) 185; Phys. Rev. Lett. **85** (2000) 3999 (SuperK) and data from Kamiokande, IMB, Soudan-2, and MACRO experiments. This data can be explained by $\nu_\mu \rightarrow \nu_\tau$ oscillations with $|\Delta m_{32}^2| \simeq 2.5 \times 10^{-3} \text{ eV}^2$ and a maximal value of the associated mixing angle factor $\sin^2 2\theta_{23}$. (The sign of Δm_{3j}^2 , $j = 1, 2$, is not determined by this data.)
 - [3] S. H. Ahn et al., Phys. Lett. **B511** (2001) 178; K. Nishikawa, talk at Neutrino-2002 (May, 2002).
 - [4] S. Weinberg, Phys. Rev. D **19** (1979) 1277; L. Susskind, Phys. Rev. D **20** (1979) 2619; E. Eichten and K. Lane, Phys. Lett. **B90** (1980) 125.
 - [5] P. Sikivie, L. Susskind, M. Voloshin, and V. Zakharov, Nuc. Phys. **B173** (1980) 189.
 - [6] B. Holdom, Phys. Lett. B **150** (1985) 301; K Yamawaki, M. Bando, and K. Matumoto, Phys. Rev. Lett. **56** (1986) 1335; T. Appelquist, D. Karabali, and L. Wijewardhana, Phys. Rev. Lett.

- 57** (1986) 957; T. Appelquist and L.C.R. Wijewardhana, Phys. Rev. D **35** (1987) 774; Phys. Rev. D **36** (1987) 568.
- [7] B. Holdom, Phys. Rev. D **23** (1981) 1637; Phys. Lett. B **246** (1990) 169.
- [8] T. Appelquist and J. Terning, Phys. Lett. **B315** (1993) 139; T. Appelquist, J. Terning, and L. Wijewardhana, Phys. Rev. Lett. **77** (1996) 1214; *ibid.* **79** (1997) 2767; T. Appelquist, N. Evans, S. Selipsky, Phys. Lett. **B374** (1996) 145; T. Appelquist and S. Selipsky, Phys. Lett. **B400** (1997) 364.
- [9] T. Appelquist, J. Terning, Phys. Rev. **D50** (1994) 2116.
- [10] Two recent reviews are R. Chivukula, hep-ph/0011264, K. Lane, hep-ph/0202255; C. Hill E. Simmons, hep-ph/0203079.
- [11] M. Gell-Mann, P. Ramond, R. Slansky, in *Supergravity* (North Holland, Amsterdam, 1979), p. 315; T. Yanagida in proceedings of *Workshop on Unified Theory and Baryon Number in the Universe*, KEK, 1979.
- [12] In general, lepton family number is violated by both Dirac and Majorana mass terms.
- [13] Although we require our model to yield small S , a reanalysis of precision electroweak data is in order in view of the recent measurement of $\sin^2 \theta_W$ by the NuTeV experiment; see G. Zeller et al., Phys. Rev. Lett. **88** (2002) 091802.
- [14] For a vectorial $SU(N)$ theory with N_f fermions in the fundamental representation, an IRFP occurs if $N_f > N_{f,min,IR}$, where, perturbatively, $N_{f,min,IR} \simeq 34N^3/(13N^2 - 3)$. At this IRFP, using the criticality condition [15], the theory is expected to exist in a confining phase with $S\chi SB$ if $N_{f,min,IR} < N_f < N_{f,con}$, where $N_{f,con} \simeq (2/5)N(50N^2 - 33)/(5N^2 - 3)$ and in a conformal phase if $N_{f,con} < N_f < 11N/2$. For $N = 2$ we have $N_{f,min,IR} \sim 5$ and $N_{f,con} \simeq 8$, respectively. For attempts at nonperturbative lattice studies of these properties, see R. Mawhinney, Nucl. Phys. Proc. Suppl. **83**, 57 (2000).
- [15] In the approximation of a single-gauge-boson exchange, the critical coupling for condensation $R_1 \times R_2 \rightarrow R_c$ is given by the condition $\frac{3\alpha}{2\pi} \Delta C_2 = 1$, where $\Delta C_2 = [C_2(R_1) + C_2(R_2) - C_2(R_c)]$, and $C_2(R)$ is the quadratic Casimir invariant. Instanton contributions are also important [8].
- [16] Here $\eta_a = \exp[\int_{f_F}^{\Lambda_a} (d\mu/\mu) \gamma(\alpha(\mu))]$, and in walking TC theories the anomalous dimension $\gamma \simeq 1$ so $\eta_a \simeq \Lambda_a/f_F$.
- [17] We write SM-singlet neutrinos as right-handed fields $\chi_{j,R}$.
- [18] Z. Maki, M. Nakagawa, S. Sakata, Prog. Theor. Phys. **28** (1962) 870 (2×2 matrix); B. W. Lee, S. Pakvasa, R. Shrock, and H. Sugawara, Phys. Rev. Lett. **38** (1977) 937 (3×3 matrix).
- [19] T. Appelquist and R. Shrock, to appear.
- [20] Since ETC is a chiral gauge theory which is strongly interacting at its self-breaking scale (here, Λ_1) and since $\pi_3(SU(N)) = \mathbb{Z}$, the associated ETC instantons will generically violate total lepton number L . Because $g_{ETC} \sim O(1)$ at Λ_1 , these instantons are not suppressed by small $e^{-16\pi^2/g^2}$ factors, in contrast to the situation at zero temperature in the weak $SU(2)$ sector. However, the resultant effective multifermion operators are of quite high dimension and are thus suppressed at low energies.
- [21] R. Shrock, Phys. Lett. **96B** (1980) 159; Phys. Rev. D **24** (1981) 1232, 1275.
- [22] Current limits are summarized in <http://pdg.lbl.gov>.